

STEP 2021, P3, Q3 - Solution (4 pages; 14/7/23)**(i) 1st Part**

$$\begin{aligned}
& \frac{1}{2}(I_{n+1} + I_{n-1}) \\
&= \frac{1}{2} \left[\int_0^\beta (\sec x + \tan x)^{n+1} dx + \int_0^\beta (\sec x + \tan x)^{n-1} dx \right] \\
&= \frac{1}{2} \int_0^\beta (\sec x + \tan x)^{n-1} (\sec^2 x + \tan^2 x + 2\sec x \tan x + 1) dx \\
&= \int_0^\beta (\sec x + \tan x)^{n-1} (\sec^2 x + \sec x \tan x) dx
\end{aligned}$$

$$\begin{aligned}
\text{Now, } \frac{d}{dx} \left[\frac{1}{n} (\sec x + \tan x)^n \right] \\
&= (\sec x + \tan x)^{n-1} [-(\cos x)^{-2}(-\sin x) + \sec^2 x] \\
&= (\sec x + \tan x)^{n-1} [\sec^2 x + \sec x \tan x]
\end{aligned}$$

$$\text{And } \frac{1}{n} (\sec(0) + \tan(0))^n = \frac{1}{n},$$

$$\begin{aligned}
\text{so that } \int_0^\beta (\sec x + \tan x)^{n-1} (\sec^2 x + \sec x \tan x) dx \\
&= \left[\frac{1}{n} (\sec x + \tan x)^n \right]_0^\beta \\
&= \frac{1}{n} (\sec \beta + \tan \beta)^n - \frac{1}{n}, \text{ as required.} \\
&= \frac{1}{n} [(\sec \beta + \tan \beta)^n - 1]
\end{aligned}$$

2nd Part**Method 1**

Suppose instead that $I_n \geq \frac{1}{n} [(\sec \beta + \tan \beta)^n - 1]$

$$\begin{aligned}
\text{Then } \frac{1}{2}(I_{n+1} + I_{n-1}) &\geq \frac{1}{2n}[(\sec\beta + \tan\beta)^{n+1} - 1] \\
&+ \frac{1}{2n}[(\sec\beta + \tan\beta)^{n-1} - 1] \\
&= \frac{1}{2n}(\sec\beta + \tan\beta)^{n-1}[(\sec\beta + \tan\beta)^2 + 1] \\
&= \frac{1}{2n}(\sec\beta + \tan\beta)^{n-1}[\sec^2\beta + \tan^2\beta + 2\sec\beta\tan\beta + 1] \\
&= \frac{1}{n}(\sec\beta + \tan\beta)^{n-1}(\sec^2\beta + \sec\beta\tan\beta) \\
&= \frac{1}{n}(\sec\beta + \tan\beta)^n \sec\beta > \frac{1}{n}(\sec\beta + \tan\beta)^n, \text{ as } 0 < \beta < \frac{\pi}{2} \quad (*)
\end{aligned}$$

But, from the 1st Part, $\frac{1}{2}(I_{n+1} + I_{n-1}) = \frac{1}{n}[(\sec\beta + \tan\beta)^n - 1]$
 $< \frac{1}{n}[(\sec\beta + \tan\beta)^n]$, which contradicts (*).

$$\text{Hence } I_n < \frac{1}{n}[(\sec\beta + \tan\beta)^n - 1]$$

Method 2

From the 1st Part, the result to prove is equivalent to

$$I_n < \frac{1}{2}(I_{n+1} + I_{n-1}) \text{ or } I_{n+1} + I_{n-1} - 2I_n > 0$$

$$\text{And } I_{n+1} + I_{n-1} - 2I_n =$$

$$\int_0^\beta (\sec x + \tan x)^{n+1} + (\sec x + \tan x)^{n-1} - 2(\sec x + \tan x)^n dx$$

$$= \int_0^\beta (\sec x + \tan x)^{n-1} (\sec^2 x + \tan^2 x + 2\sec x \tan x + 1$$

$$- 2\sec x - 2\tan x) dx$$

$$= 2 \int_0^\beta (\sec x + \tan x)^{n-1} (\sec^2 x + \sec x \tan x - \sec x - \tan x) dx$$

$$= 2 \int_0^\beta (\sec x + \tan x)^{n-1} (\sec x + \tan x)(\sec x - 1) dx$$

$$= 2 \int_0^\beta (\sec x + \tan x)^n (\sec x - 1) dx > 0, \text{ as required,}$$

as both $\sec x + \tan x$ and $\sec x - 1$ are positive for $0 < x < \beta < \frac{\pi}{2}$

(ii) [It isn't clear what approach the question setter has in mind here. Possible options are:

(a) Applying exactly the same method; ie starting by showing that

$$\frac{1}{2}(J_{n+1} + J_{n-1}) = \frac{1}{n}((1 + \tan x)^n - \cos^n x) \text{ [This gives the}$$

following for the LHS:

$$\frac{1}{2} \left[\int_0^\beta (\sec x \cos \beta + \tan x)^{n+1} dx + \int_0^\beta (\sec x \cos \beta + \tan x)^{n-1} dx \right]$$

$$= \frac{1}{2} \int_0^\beta (\sec x \cos \beta + \tan x)^{n-1} (\sec^2 x \cos^2 \beta + \tan^2 x$$

$+ 2 \sec x \sin x + 1) dx$, which isn't very promising.]

(b) Modifying the method in some way that takes account of the differences between J_n and I_n . Nothing obvious springs to mind.

(c) Using the result of Part (i) in some way; eg by making a substitution. Again, nothing obvious springs to mind.

(d) Using an idea that was involved in answering Part (i).

One idea was that of showing that an integral had a positive value. In order to use this we will need to be able to write

$$\frac{1}{n}((1 + \tan x)^n - \cos^n x) \text{ as in integral } (K_n, \text{ say}), \text{ and show that}$$

$$K_n - J_n > 0$$

Another idea was using the fact that certain components of the integrand were positive.

(e) Using an idea prompted by a difference between J_n and I_n .

One such idea that emerges later on is that $\cos\beta < \cos x$ for $0 < x < \beta$, and this enables the awkward $\sec x \cos\beta$ to be converted into $\sec x \cos x = 1$]

$$\text{Consider } \frac{d}{dx} \left[\frac{1}{n} ((1 + \tan x)^n - \cos^n x) \right]$$

$$= (1 + \tan x)^{n-1} \sec^2 x - \cos^{n-1} x (-\sin x)$$

$$\text{Let } K_n = \int_0^\beta (1 + \tan x)^{n-1} \sec^2 x + \cos^{n-1} x \sin x dx$$

$$= \left[\frac{1}{n} ((1 + \tan x)^n - \cos^n x) \right]_0^\beta$$

$$= \frac{1}{n} ((1 + \tan\beta)^n - \cos^n \beta) - 0$$

Then the result to be proved is that $K_n - J_n > 0$

$$\text{Now, } K_n - J_n = \int_0^\beta (1 + \tan x)^{n-1} \sec^2 x + \cos^{n-1} x \sin x$$

$$- (\sec x \cos\beta + \tan x)^{n+1} dx$$

$$> \int_0^\beta (1 + \tan x)^{n-1} \sec^2 x + \cos^{n-1} x \sin x$$

$$- (\sec x \cos x + \tan x)^{n+1} dx,$$

$$\text{as } x < \beta \Rightarrow \cos x > \cos\beta \text{ (for } 0 < \beta < \frac{\pi}{2}),$$

$$= \int_0^\beta (1 + \tan x)^{n-1} \sec^2 x + \cos^{n-1} x \sin x - (1 + \tan x)^{n+1} dx,$$

$$= \int_0^\beta (1 + \tan x)^{n-1} (\tan^2 x + 1) + \cos^{n-1} x \sin x$$

$$- (1 + \tan x)^{n+1} dx$$

$$= \int_0^\beta (1 + \tan x)^{n-1} + \cos^{n-1} x \sin x > 0, \text{ as required,}$$

as $1 + \tan x$, $\cos x$ & $\sin x$ are all positive for $0 < x < \beta < \frac{\pi}{2}$