


## Hooke's Law - Exercises (Solutions)

(5 pages; 26/1/19)

(1) A particle of mass 200g is attached at the mid-point of an elastic string of natural length 0.5m and modulus of elasticity  $\lambda$ , which hangs vertically between two points, 1m apart.

- (i) How far will the particle be below the top point if  $\lambda = 1$ ?
- (ii) Determine the minimum value of  $\lambda$  such that there is no slack in the string.

### Solution



(i) Let the extensions of the upper and lower parts of the string be  $e_u$  and  $e_l$ , respectively, and the tensions in the two parts  $T_u$  and  $T_l$ .

Then, referring to the diagram,

$$T_u = \frac{\lambda e_u}{0.25} \quad , \quad T_l = \frac{\lambda e_l}{0.25} \quad (\text{assuming the string is not slack})$$

$$\text{Equilibrium} \Rightarrow T_u = T_l + 0.2g \quad ; \text{ also } e_u + e_l = 0.5 \quad (1)$$

$$\text{Hence } \lambda e_u = \lambda e_l + 0.05g$$

$$\text{and so } \lambda e_u = \lambda(0.5 - e_u) + 0.05g,$$

$$\text{giving } 2\lambda e_u = 0.5\lambda + 0.05g$$

$$\text{and hence } e_u = \frac{0.5\lambda + 0.05g}{2\lambda} \quad (2)$$

$$\text{Thus when } \lambda = 1, e_u = 0.495$$

$$\text{and the distance below the top point is } 0.25 + 0.495 = 0.745m$$

(ii) The string is slack if  $e_l < 0$

$$\text{From (1) \& (2), } e_l = 0.5 - 0.25 - \frac{0.025g}{\lambda} = 0.25 - \frac{0.025g}{\lambda}$$

$$\text{Thus we require } 0.25 - \frac{0.025g}{\lambda} \geq 0,$$

$$\text{so that } 0.25 \geq \frac{0.025g}{\lambda} \quad \text{and } \lambda \geq 0.1g = 0.98$$


(2) (Oscillations)

A particle of mass 200g hangs at a point Q, suspended from a fixed point P, by means of a spring of original length 20cm and modulus of elasticity 5N. It is pulled down to a point R, which is 35cm below P. The particle is then released.

Ignoring any resistances to motion, find:

- (i) the work done in pulling the particle down to R
- (ii) the maximum speed of the particle after it is released, and the point at which this occurs
- (iii) the distance of the particle below P when it reaches its maximum height, at position S, and show that the distance QS equals the distance QR

### Solution



[Note: The  $g$  in the diagram (gravity) is not to be confused with  $g$  for grams.]

- (i) If  $e$  is the extension of the spring at Q (in metres),

$$\text{Hooke's Law} \Rightarrow \frac{\lambda e}{l} = T = mg \Rightarrow \frac{5e}{0.2} = (0.2)(9.8) \Rightarrow e = 0.0784$$

Taking the zero of gravitational potential energy (GPE) to be R, the total energy of the particle at Q is:

GPE + EPE + KE (where EPE is elastic potential energy & KE is kinetic energy)

$$= (0.2)(9.8)(0.35 - 0.2 - 0.0784) + \frac{1}{2} \left( \frac{5}{0.2} \right) (0.0784)^2 + 0$$

$$= 0.140336 + 0.076832 + 0 = 0.217168$$

The total energy of the particle at R is:

$$0 + \frac{1}{2} \left( \frac{5}{0.2} \right) (0.15)^2 + 0 = 0.28125$$

$$\text{Thus the work done} = 0.28125 - 0.217168 = 0.064082 = 0.0641 \text{ J (3sf)}$$

(ii) The maximum speed will occur when the particle is not accelerating; ie at Q, where the net force on the particle is zero [as  $T = mg$  at the equilibrium position].

The KE of the particle at Q will equal the work done to pull it down to R, as this is the energy gained by the particle since it was last at Q.

$$\text{Hence } \frac{1}{2} (0.2)v^2 = 0.064082 \text{ (where } v \text{ is the maximum speed)}$$

$$\text{and } v = 0.80051 = 0.801 \text{ ms}^{-1} \text{ (3sf)}$$

(iii) Let  $d$  be the distance below P when the particle is at S.

The total energy of the particle at S is:

$$(0.2)(9.8)(0.35 - d) + \frac{1}{2} \left( \frac{5}{0.2} \right) (d - 0.2)^2 + 0$$

and this equals the energy at R of 0.28125, so that


$$12.5d^2 - 6.96d + 0.90475 = 0$$

$$\text{and } d = \frac{6.96 \pm \sqrt{3.2041}}{25} = 0.35 \text{ or } 0.2068$$

Thus 0.35 corresponds to R and S is the point 20.68cm below P.

This is  $20 + 7.84 - 20.68 = 7.16$  cm above the equilibrium position Q, whilst R is  $35 - (20 + 7.84) = 7.16$  cm below Q.

[The particle oscillates between R and S.]



### (3) (Elastic Potential Energy)

A bungee jumper of mass 80kg is attached to a rope of original length 10m and modulus of elasticity 1600N. How far will he or she fall? (Take  $g=10$ )

#### Solution

Let  $e$  be the extension of the rope.

Gain in elastic PE = loss of gravitational PE, so that

$$\frac{1}{2} \left( \frac{1600}{10} \right) e^2 = 80(10)(10 + e)$$

$$\Rightarrow e^2 = 100 + 10e$$

$$\Rightarrow e^2 - 10e - 100 = 0$$

$$\Rightarrow e = \frac{10 \pm \sqrt{100 + 400}}{2} = 16.18\text{m (ignoring -ve value)}$$

So bungee jumper falls by  $10 + 16.18 = 26.18\text{m}$