

Hyperbolas - Exercises (Solutions) (6 pages; 10/9/19)

(1) Show that the equation of the tangent to the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \text{ at the point } (a\cosh t, b\sinh t) \text{ is}$$

$$y\sinh t = x\cosh t - ab$$

Solution

Using the parametric equations $x = a\cosh t$ & $y = b\sinh t$,

$$\frac{dx}{dt} = a\sinh t \quad \& \quad \frac{dy}{dt} = b\cosh t,$$

$$\text{so that } \frac{dy}{dx} = \frac{b\cosh t}{a\sinh t}$$

and the equation of the tangent at $(a\cosh t, b\sinh t)$ is

$$\frac{y - b\sinh t}{x - a\cosh t} = \frac{b\cosh t}{a\sinh t}$$

$$\text{and hence } y\sinh t - ab\sinh^2 t = x\cosh t - ab\cosh^2 t,$$

$$\text{so that } y\sinh t = x\cosh t - ab$$

(2) Given that the tangent in (C)(1) meets the asymptotes of the hyperbola at the points P & Q , show that the mid-point of P & Q is $(a\cosh t, b\sinh t)$.

Solution

The asymptotes of the hyperbola are $y = \pm \frac{b}{a}x$

From (1), the tangent to the hyperbola at $(a\cosh t, b\sinh t)$ meets the asymptote $y = \frac{b}{a}x$ at P (say), where $b\sinh t = x\cosh t - ab$ and the asymptote $y = -\frac{b}{a}x$ at Q where

$$-b\sinh t = x\cosh t - ab$$

so that P is the point $\left(\frac{a}{\cosh t - \sinh t}, \frac{b}{\cosh t - \sinh t}\right)$

and Q is the point $\left(\frac{a}{\cosh t + \sinh t}, \frac{-b}{\cosh t + \sinh t}\right)$

The mid-point of P & Q is therefore

$$\begin{aligned} & \left(\frac{1}{2} \left[\frac{a}{\cosh t - \sinh t} + \frac{a}{\cosh t + \sinh t}\right], \frac{1}{2} \left[\frac{b}{\cosh t - \sinh t} + \frac{-b}{\cosh t + \sinh t}\right]\right) \\ &= \left(\frac{a \cosh t}{\cosh^2 t - \sinh^2 t}, \frac{b \sinh t}{\cosh^2 t - \sinh^2 t}\right) = (a \cosh t, b \sinh t), \text{ as required.} \end{aligned}$$

(3) In the case where $b = a$, find the area of the triangle OPQ (where P & Q are as in (C)(2), and O is the Origin).

Solution

The two asymptotes are at right angles to each other, so that the required area, $A = \frac{1}{2} OP \cdot OQ$

$$\begin{aligned} \text{Then } 4A^2 &= \left(\left(\frac{a}{\cosh t - \sinh t}\right)^2 + \left(\frac{a}{\cosh t - \sinh t}\right)^2\right) \\ &\times \left(\left(\frac{a}{\cosh t + \sinh t}\right)^2 + \left(\frac{-a}{\cosh t + \sinh t}\right)^2\right) \\ &= \left(\frac{2a^2}{(\cosh t - \sinh t)^2}\right) \left(\frac{2a^2}{(\cosh t + \sinh t)^2}\right) \\ &= \frac{4a^4}{(\cosh^2 t - \sinh^2 t)^2} = 4a^4 \end{aligned}$$

and therefore $A = a^2$

(4) The chord PQ , where P and Q are points on the rectangular hyperbola $xy = c^2$, has gradient 1. Show that the locus of the point of intersection of the tangents from P and Q is the line

$$y = -x. \text{ [Edx FP3 textbook, Ex. 2G, Q9]}$$

Solution

Let P & Q be the points $(ct_1, \frac{c}{t_1})$ & $(ct_2, \frac{c}{t_2})$, respectively.

As the gradient of PQ is 1, $\frac{\frac{c}{t_2} - \frac{c}{t_1}}{ct_2 - ct_1} = 1$, so that

$$\frac{1}{t_2} - \frac{1}{t_1} = t_2 - t_1$$

$$\Rightarrow \frac{t_1 - t_2}{t_1 t_2} = t_2 - t_1$$

$$\Rightarrow t_1 t_2 = -1, \text{ as } t_1 \neq t_2 \text{ (} P \text{ \& } Q \text{ being distinct points)}$$

The equation of the tangent from P is:

$$\frac{y - \frac{c}{t_1}}{x - ct_1} = \frac{dy/dt}{dx/dt} \Big|_{t=t_1}, \text{ where } x = ct \text{ \& } y = \frac{c}{t}$$

$$\text{so that } \frac{dy}{dt} = -\frac{c}{t^2} \text{ \& } \frac{dx}{dt} = c$$

and the equation of the tangent from P is

$$\frac{y - \frac{c}{t_1}}{x - ct_1} = \left(\frac{-\frac{c}{t_1^2}}{c} \right) \Rightarrow t_1^2 y - t_1 c = -(x - ct_1)$$

$$\Rightarrow t_1^2 y = -x + 2ct_1 \quad (1)$$

Similarly, the equation of the tangent from Q is $t_2^2 y = -x + 2ct_2$

and these lines intersect where

$$t_1^2 y - 2ct_1 = t_2^2 y - 2ct_2,$$

$$\text{so that } y(t_1^2 - t_2^2) = 2c(t_1 - t_2)$$

$$\text{and } y = \frac{2c}{t_1 + t_2} \text{ (as } t_1 \neq t_2)$$

$$\text{Then, from (1), } x = 2ct_1 - \frac{2ct_1^2}{t_1 + t_2}$$

$$= \frac{2ct_1^2 + 2ct_1t_2 - 2ct_1^2}{t_1 + t_2}$$

$$= \frac{2ct_1t_2}{t_1 + t_2}$$

and so $\frac{y}{x} = \frac{1}{t_1t_2} = -1$ (found earlier),

and thus the points of intersection satisfy $y = -x$, as required.

(5) Use matrices to show that the rectangular hyperbola $x^2 - y^2 = a^2$ can be obtained by rotating the rectangular hyperbola $xy = c^2$, expressing a^2 in terms of c .

Solution

The asymptotes of $x^2 - y^2 = a^2$ are $y = \pm x$, whilst the asymptotes of

$xy = c^2$ are the x and y axes.

So consider a rotation of 45° clockwise.

Then the point (x, y) on the hyperbola $xy = c^2$ is transformed to the point (u, v) , where

$$\begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} \cos(-45^\circ) & -\sin(-45^\circ) \\ \sin(-45^\circ) & \cos(-45^\circ) \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

$$= \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

$$= \frac{1}{\sqrt{2}} \begin{pmatrix} x + y \\ -x + y \end{pmatrix}$$

Then $u^2 - v^2 = (u - v)(u + v)$

$$= \frac{1}{\sqrt{2}}(2x) \cdot \frac{1}{\sqrt{2}}(2y) = 2xy = 2c^2$$

Relabelling gives $x^2 - y^2 = 2c^2$ (and $a^2 = 2c^2$).

(6) Show that the equation of the normal to the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \text{ at the point } (a\cosh t, b\sinh t) \text{ is}$$

$$x\sinh t + y\cosh t = (a^2 + b^2)\sinh t\cosh t$$

Solution

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{b\cosh t}{a\sinh t}$$

$$\text{so that equation of normal is } y - b\sinh t = -\frac{a\sinh t}{b\cosh t}(x - a\cosh t)$$

$$\Rightarrow b\cosh t \cdot y - b^2\sinh t\cosh t = -x\sinh t + a^2\sinh t\cosh t$$

$$\Rightarrow x\sinh t + y\cosh t = (a^2 + b^2)\sinh t\cosh t, \text{ as required}$$

(7) [Edx (Pearson), FP1, ME2, Q7 (p58)]

l_1 & l_2 are distinct tangents to the rectangular hyperbola $xy = 9$ with gradient $-\frac{1}{4}$; find the equations of l_1 & l_2

Solution

$$y = \frac{9}{x} \Rightarrow \frac{dy}{dx} = -9x^{-2}$$

Suppose that the gradient at the point $(a, \frac{9}{a})$ is $-\frac{1}{4}$

Then $-9a^{-2} = -\frac{1}{4}$, so that $a^2 = 36$, and $a = \pm 6$

$$\text{Equation of } l_1: y - \frac{9}{6} = -\frac{1}{4}(x - 6);$$

$$\text{or } 4y - 6 = -x + 6; x + 4y = 12$$

$$\text{Equation of } l_2: y - (-\frac{9}{6}) = -\frac{1}{4}(x - [-6]);$$

$$\text{or } 4y + 6 = -x - 6; x + 4y = -12$$

(8) [Edx (Pearson), FP1, ME2, Q14 (p59)]

Suppose that P is a general point on a rectangular hyperbola and that the tangent at P crosses the x and y axes at A and B respectively. Show that:

(i) $AP = BP$

(ii) the triangle OAB has a constant area, as P varies

Solution

(i) Suppose that the equation of the rectangular hyperbola is $xy = c^2$ and that P is the point $\left(a, \frac{c^2}{a}\right)$.

Then $\frac{dy}{dx} = -c^2x^{-2}$, and the equation of the tangent at P is

$$y - \frac{c^2}{a} = -\frac{c^2}{a^2}(x - a)$$

At A, $0 - \frac{c^2}{a} = -\frac{c^2}{a^2}(x - a)$, so that $a = x - a$ and $x = 2a$

At B, $y - \frac{c^2}{a} = -\frac{c^2}{a^2}(0 - a)$, so that $y = \frac{c^2}{a} + \frac{c^2}{a} = \frac{2c^2}{a}$

$$\text{Then } AP^2 = (2a - a)^2 + \left(\frac{c^2}{a} - 0\right)^2 = a^2 + \frac{c^4}{a^2}$$

$$\text{and } BP^2 = (a - 0)^2 + \left(\frac{2c^2}{a} - \frac{c^2}{a}\right)^2 = a^2 + \frac{c^4}{a^2}$$

Thus $AP = BP$, as required.

(ii) Area of OAB = $\frac{1}{2}(2a)\left(\frac{2c^2}{a}\right) = 2c^2$; ie a constant