

## Hyperbolic Functions: Exercises - Sol'ns (13 pages; 28/9/19)

(1) (i) Prove, using exponential functions, that

(a)  $\cosh^2 x - \sinh^2 x = 1$

(b)  $\sinh 2x = 2 \sinh x \cosh x$

(ii) By differentiating the result from (i)(b), obtain an expression for  $\cosh 2x$  in terms of  $\cosh^2 x$  and  $\sinh^2 x$

### Solution

(i)(a) As  $\cosh x = \frac{1}{2}(e^x + e^{-x})$  &  $\sinh x = \frac{1}{2}(e^x - e^{-x})$ ,

$$\begin{aligned} \cosh^2 x - \sinh^2 x &= (\cosh x + \sinh x)(\cosh x - \sinh x) \\ &= e^x \cdot e^{-x} = 1 \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad 2 \sinh x \cosh x &= 2 \left(\frac{1}{2}\right)(e^x - e^{-x}) \left(\frac{1}{2}\right)(e^x + e^{-x}) \\ &= \frac{1}{2}(e^{2x} - e^{-2x}) = \sinh 2x \quad (\text{by difference of 2 squares}) \end{aligned}$$

(ii) Differentiating  $\sinh 2x = 2 \sinh x \cosh x$  gives

$$\begin{aligned} 2 \cosh 2x &= 2 \cosh x \cosh x + 2 \sinh x \sinh x \\ \Rightarrow \cosh 2x &= \cosh^2 x + \sinh^2 x \end{aligned}$$

(2) (a) Find the formula connecting  $\tanh^2 x$  &  $\operatorname{sech}^2 x$ ?

(b) Find the formula connecting  $\operatorname{coth}^2 x$  &  $\operatorname{cosech}^2 x$ ?

### Solution

From  $\cosh^2 x - \sinh^2 x = 1$ ,

(a) divide by  $\cosh^2 x$ , to give  $1 - \tanh^2 x = \operatorname{sech}^2 x$

(b) divide by  $\sinh^2 x$ , to give  $\operatorname{coth}^2 x - 1 = \operatorname{cosech}^2 x$

(3) Show that  $\operatorname{artanh} x = \frac{1}{2} \ln \left( \frac{1+x}{1-x} \right)$  ( $|x| < 1$ )

### Solution

If  $y = \operatorname{artanh} x$ , then  $\operatorname{tanh} y = x$  ( $|x| < 1$ )

$$\Rightarrow x = \frac{\frac{1}{2}(e^y - e^{-y})}{\frac{1}{2}(e^y + e^{-y})} = \frac{e^{2y} - 1}{e^{2y} + 1}$$

$$\Rightarrow x(e^{2y} + 1) = e^{2y} - 1$$

$$\Rightarrow e^{2y}(x - 1) = -1 - x$$

$$\Rightarrow e^{2y} = \frac{1+x}{1-x}$$

$$\Rightarrow y = \frac{1}{2} \ln \left( \frac{1+x}{1-x} \right) \quad (|x| < 1)$$

### (4) Differentiation

Find or prove the following:

(i)  $\frac{d}{dx} \operatorname{tanh} x$

(ii)  $\frac{d}{dx} \operatorname{arcosh} x = \frac{1}{\sqrt{x^2 - 1}}$

(iii)  $\frac{d}{dx} \operatorname{artanh} x = \frac{1}{1-x^2}$

(iv)  $\frac{d}{dx} \operatorname{sech} x$

### Solutions

$$\begin{aligned} \text{(i)} \quad \frac{d}{dx} \operatorname{tanh} x &= \frac{d}{dx} \frac{\sinh x}{\cosh x} = \frac{\cosh x \cdot \cosh x - \sinh x \cdot \sinh x}{\cosh^2 x} \\ &= \frac{1}{\cosh^2 x} = \operatorname{sech}^2 x \end{aligned}$$

(ii) Let  $y = \operatorname{arcosh} x$ , so that  $\operatorname{cosh} y = x$

Then  $\frac{dx}{dy} = \sinh y$  and  $\frac{dy}{dx} = \frac{1}{\sqrt{\cosh^2 y - 1}} = \frac{1}{\sqrt{x^2 - 1}}$

(iii) Let  $y = \operatorname{artanh} x$ , so that  $\tanh y = x$

and  $\frac{dx}{dy} = \operatorname{sech}^2 y = 1 - \tanh^2 y = 1 - x^2$

Hence  $\frac{d}{dx} \operatorname{artanh} x = \frac{1}{1-x^2}$

(iv)  $\frac{d}{dx} \operatorname{sech} x = \frac{d}{dx} (\cosh x)^{-1} = (-1)(\cosh x)^{-2} \sinh x$

$= -\operatorname{sech}^2 x \cdot \sinh x$  or  $-\operatorname{sech} x \cdot \tanh x$

[Although this is similar to  $\frac{d}{dx} \sec x = \sec x \cdot \tan x$ , Osborn's rule doesn't apply to derivatives (and, in any case, there is no  $\sinh^2 x$  or similar term).]

(5) Simplify  $\sinh(\cosh^{-1} 2)$

**Solution**

Let  $\cosh^{-1} 2 = a (> 0)$ , so that  $2 = \cosh a$

Then  $\sinh a = +\sqrt{\cosh^2 a - 1}$  [as  $a > 0$ ]  $= \sqrt{3}$

(6) Solve the equation  $5\cosh 2x + 3\sinh x = 6$ ,

giving your answers in exact logarithmic form

**Solution**

$$5\cosh 2x + 3\sinh x = 6 \Rightarrow 5(\cosh^2 x + \sinh^2 x) + 3\sinh x - 6 = 0$$

$$\Rightarrow 5(1 + 2\sinh^2 x) + 3\sinh x - 6 = 0$$

$$\Rightarrow 10\sinh^2 x + 3\sinh x - 1 = 0$$

$$\Rightarrow (5\sinh x - 1)(2\sinh x + 1) = 0$$

$$\Rightarrow \sinh x = \frac{1}{5} \text{ or } -\frac{1}{2}$$

$$\Rightarrow x = \operatorname{arsinh}\left(\frac{1}{5}\right) \text{ or } \operatorname{arsinh}\left(-\frac{1}{2}\right)$$

$$\Rightarrow x = \ln\left(\frac{1}{5} + \sqrt{\frac{1}{25} + 1}\right) \text{ or } \ln\left(-\frac{1}{2} + \sqrt{\frac{1}{4} + 1}\right)$$

$$\text{or } \ln\left(\frac{1}{5}(1 + \sqrt{26})\right) \text{ or } \ln\left(\frac{1}{2}(\sqrt{5} - 1)\right)$$

[It is possible to substitute these values into the equation, as a check.]

$$(7) \text{ Show that } \operatorname{arcosh} x = \ln(x + \sqrt{x^2 - 1})$$

### Solution

If  $y = \operatorname{arcosh} x$ , then  $\cosh y = x$

$$\Rightarrow x = \frac{1}{2}(e^y + e^{-y})$$

$$\Rightarrow 2xe^y = e^{2y} + 1$$

$$\Rightarrow e^{2y} - 2xe^y + 1 = 0$$

$$\Rightarrow e^y = \frac{2x \pm \sqrt{4x^2 - 4}}{2} = x \pm \sqrt{x^2 - 1}$$

$$\Rightarrow y = \ln(x \pm \sqrt{x^2 - 1})$$

However, in order for  $y = \operatorname{arcosh} x$  to be a function, the negative branch is suppressed (by restricting the domain of  $\cosh x$  to non-negative values). And we can show that  $\ln(x - \sqrt{x^2 - 1}) < 0$ :

### Method 1

Equivalently, we need to show that  $x - \sqrt{x^2 - 1} < 1$ ; or that

$$x - 1 < \sqrt{x^2 - 1}$$

But  $x - 1 = \sqrt{(x - 1)(x - 1)}$  (noting that the range of  $\cosh x$ , and hence the domain of  $\operatorname{arcosh} x$ , excludes  $x < 1$ )

and  $\sqrt{(x - 1)(x - 1)} < \sqrt{(x - 1)(x + 1)} = \sqrt{x^2 - 1}$ , as required

( $y = \sqrt{x}$  is an increasing function,

so  $x - 1 < x + 1 \Rightarrow \sqrt{x - 1} < \sqrt{x + 1}$  )

[Alternatively, we can argue (slightly informally) that the difference between  $x^2$  and  $x^2 - 1$  (ie 1) is contracted by applying the square root function, so that  $\sqrt{x^2} - \sqrt{x^2 - 1} < 1$ ]

## Method 2

We expect the unrestricted  $y = \operatorname{arcosh} x$  to be symmetric about the  $x$ -axis (as  $y = \cosh x$  is symmetric about the  $y$ -axis). So we could show that  $y = \ln(x \pm \sqrt{x^2 - 1})$  can also be written as

$y = \pm \ln(x + \sqrt{x^2 - 1})$ , and then reject the negative branch as before.

So we want to show that  $\ln(x - \sqrt{x^2 - 1}) = -\ln(x + \sqrt{x^2 - 1})$ :

$$\text{RHS} = \ln\left(\frac{1}{x + \sqrt{x^2 - 1}}\right) = \ln\left(\frac{x - \sqrt{x^2 - 1}}{x^2 - (x^2 - 1)}\right) = \ln(x - \sqrt{x^2 - 1}), \text{ as}$$

required

(8) If  $x = \sinh u$ , write  $\sinh(4u)$  in terms of  $x$

## Solution

$$\begin{aligned} \sinh(4u) &= 2 \sinh(2u) \cosh(2u) \\ &= 4 \sinh u \cosh u (\cosh^2 u + \sinh^2 u) \\ &= 4x \sqrt{1 + x^2} (1 + 2x^2) \end{aligned}$$

(9) Derive an expression for  $\operatorname{arsinh}(a)$  in the form  $\ln b$

**Solution**

Let  $x = \operatorname{arsinh}(a)$ , so that  $\sinh x = a$

$$\text{and } \frac{1}{2}(e^x - e^{-x}) = a$$

$$\text{Then } \frac{1}{2}(e^{2x} - 1) = ae^x$$

$$\text{and } e^{2x} - 2ae^x - 1 = 0,$$

so that  $e^x = \frac{2a \pm \sqrt{4a^2 + 4}}{2} = a + \sqrt{a^2 + 1}$  (rejecting the negative root)

$$\text{Thus } \operatorname{arsinh}(a) = \ln(a + \sqrt{a^2 + 1})$$

(noting that  $a + \sqrt{a^2 + 1} > 0$ )

$$\text{Note that } \operatorname{arsinh}\left(\frac{x}{a}\right) = \ln\left(\frac{x}{a} + \sqrt{\left(\frac{x}{a}\right)^2 + 1}\right)$$

$$= \ln\left(\frac{x + \sqrt{x^2 + a^2}}{a}\right) = \ln(x + \sqrt{x^2 + a^2}) - \ln a$$

In the formulae booklets,  $\int \frac{1}{\sqrt{a^2 + x^2}} dx$  is often given as

" $\operatorname{arsinh}\left(\frac{x}{a}\right)$  or  $\ln(x + \sqrt{x^2 + a^2})$ " but, as we've just seen, these two expressions differ by a constant

(10) Given that  $\operatorname{artanh} x = \frac{1}{2} \ln\left(\frac{1+x}{1-x}\right)$  and  $\operatorname{arcoth} x = \frac{1}{2} \ln\left(\frac{1+x}{x-1}\right)$ ,

$$\text{and also that } \frac{d}{dx}(\operatorname{artanh} x) = \frac{d}{dx}(\operatorname{arcoth} x) = \frac{1}{1-x^2},$$

what is wrong with the following reasoning?

$$\int \frac{1}{1-x^2} dx = \operatorname{artanh} x + C = \operatorname{arcoth} x + C_1,$$

so that  $\operatorname{artanh}x - \operatorname{arcoth}x = C_2$

$$\text{But } \operatorname{artanh}x - \operatorname{arcoth}x = \frac{1}{2} \ln \left( \frac{\left(\frac{1+x}{1-x}\right)}{\left(\frac{1+x}{x-1}\right)} \right) = \frac{1}{2} \ln \left( \frac{x-1}{1-x} \right) = \frac{1}{2} \ln(-1),$$



which isn't defined!

### Solution

The problem is that the domains of  $y = \operatorname{artanh}x$  and

$y = \operatorname{arcoth}x$  don't overlap (see graphs below). We ought to say that  $\operatorname{artanh}x = \frac{1}{2} \ln \left( \frac{1+x}{1-x} \right)$  for  $|x| < 1$  and  $\operatorname{arcoth}x = \frac{1}{2} \ln \left( \frac{1+x}{x-1} \right)$  for  $|x| > 1$ . So it doesn't make sense to determine

$\operatorname{artanh}x - \operatorname{arcoth}x$



Note that, with  $|x| < 1$ ,  $\frac{d}{dx}(\operatorname{artanh}x) = \frac{1}{1-x^2} > 0$  for all  $x$ ; whilst


with  $|x| > 1$ ,  $\frac{d}{dx}(\operatorname{arcoth}x) = \frac{1}{1-x^2} < 0$  for all  $x$

(11) Given that  $\sinh x = \tan y$ , where  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ , show that

(a)  $\tanh x = \sin y$  (b)  $x = \ln(\tan y + \sec y)$

### Solution

(a) As  $\sinh x = \tan y$ , we can construct a right-angled triangle (see diagram below), where the hypotenuse is  $\cosh x$ , as  $\sinh^2 x + 1 = \cosh^2 x$ .



Then  $\sin y = \frac{\sinh x}{\cosh x} = \tanh x$ , as required.

Alternatively:  $\tanh x = \frac{\sinh x}{\cosh x} = \frac{\tan y}{\sqrt{1 + \sinh^2 x}}$

(from  $\sinh^2 x + 1 = \cosh^2 x$ , noting that  $\cosh x$  is always positive, so that we take the positive square root)

$$= \frac{\tan y}{\sqrt{1 + \tan^2 y}} = \frac{\tan y}{\sqrt{\sec^2 y}} = \frac{\tan y}{\sec y}$$

(as  $\cos y > 0$  when  $-\frac{\pi}{2} < y < \frac{\pi}{2}$ , and hence  $\sec y > 0$  also)

$$= \tan y \cos y = \sin y$$

(b) From the right-angled triangle,

$$\tan y + \sec y = \sinh x + \cosh x$$

$$= \frac{1}{2}(e^x - e^{-x}) + \frac{1}{2}(e^x + e^{-x}) = e^x,$$



so that  $\ln(\tanh y + \operatorname{sech} y) = x$ , as required.

**Alternatively:**  $\sinh x = \tanh y \Rightarrow \frac{1}{2}(e^x - e^{-x}) = \tanh y$

$$\Rightarrow e^{2x} - 1 = 2 \tanh y e^x$$

$$\Rightarrow e^{2x} - 2 \tanh y e^x - 1 = 0$$

$$\Rightarrow e^x = \frac{2 \tanh y \pm \sqrt{4 \tanh^2 y + 4}}{2} = \tanh y \pm \operatorname{sech} y$$

$$\tanh y - \operatorname{sech} y = \frac{\sinh y - 1}{\cosh y} < 0 \text{ when } -\frac{\pi}{2} < y < \frac{\pi}{2}$$

Hence, as  $e^x > 0$ , it follows that  $e^x = \tanh y + \operatorname{sech} y$ ,

and hence  $x = \ln(\tanh y + \operatorname{sech} y)$

(12) What is the domain of  $\operatorname{artanh}\left(\frac{x}{2}\right)$ ?

**Solution**

$$\tanh x = \frac{\sinh x}{\cosh x} = \frac{e^x - e^{-x}}{e^x + e^{-x}} = \frac{e^{2x} - 1}{e^{2x} + 1} = \frac{e^{2x} + 1}{e^{2x} + 1} - \frac{2}{e^{2x} + 1} = 1 - \frac{2}{e^{2x} + 1}$$

Thus  $-1 < \tanh x < 1$  (as  $x \rightarrow -\infty$  &  $\infty$ )

As  $\operatorname{artanh} x$  is the inverse of  $\tanh x$ , the domain of  $\operatorname{artanh} x$  is the range of  $\tanh x$ ; ie  $(-1, 1)$ .

Thus the domain of  $\operatorname{artanh}\left(\frac{x}{2}\right)$  satisfies  $-1 < \frac{x}{2} < 1$ ;

ie  $-2 < x < 2$

(13) Show that  $\operatorname{arcoth} x = \frac{1}{2} \ln\left(\frac{1+x}{x-1}\right)$  ( $|x| > 1$ )

**Solution**

If  $y = \operatorname{arcoth} x$ , then  $\operatorname{coth} y = x$  ( $|x| > 1$ )

$$\begin{aligned} \Rightarrow x &= \frac{\frac{1}{2}(e^y + e^{-y})}{\frac{1}{2}(e^y - e^{-y})} = \frac{e^{2y} + 1}{e^{2y} - 1} \\ \Rightarrow x(e^{2y} - 1) &= e^{2y} + 1 \\ \Rightarrow e^{2y}(x - 1) &= 1 + x \\ \Rightarrow e^{2y} &= \frac{1+x}{x-1} \\ \Rightarrow y &= \frac{1}{2} \ln \left( \frac{1+x}{x-1} \right) \quad (|x| > 1) \end{aligned}$$

### Alternative Method

If  $\operatorname{artanh} x = \frac{1}{2} \ln \left( \frac{1+x}{1-x} \right)$  ( $|x| < 1$ ) has been established:

If  $y = \operatorname{arcoth} x$ , then  $\operatorname{coth} y = x$

$$\Rightarrow \operatorname{tanh} y = \frac{1}{x},$$

and hence  $y = \operatorname{artanh} \left( \frac{1}{x} \right) = \frac{1}{2} \ln \left( \frac{1+\frac{1}{x}}{1-\frac{1}{x}} \right) = \frac{1}{2} \ln \left( \frac{x+1}{x-1} \right)$

(14) (i) Use  $\operatorname{artanh} x = \frac{1}{2} \ln \left( \frac{1+x}{1-x} \right)$  to show that  $\frac{d}{dx} \operatorname{artanh} x = \frac{1}{1-x^2}$

(ii) Use  $\operatorname{arcoth} x = \frac{1}{2} \ln \left( \frac{1+x}{x-1} \right)$  to show that  $\frac{d}{dx} \operatorname{arcoth} x = \frac{1}{1-x^2}$  also

### Solution

$$\begin{aligned} \text{(i)} \quad \frac{d}{dx} \operatorname{artanh} x &= \frac{1}{2} \cdot \frac{1-x}{1+x} \cdot \frac{(1-x) - (1+x)(-1)}{(1-x)^2} \\ &= \frac{1}{2} \cdot \frac{2}{(1+x)(1-x)} = \frac{1}{1-x^2} \end{aligned}$$

$$\text{(ii)} \quad \frac{d}{dx} \operatorname{arcoth} x = \frac{1}{2} \cdot \frac{x-1}{1+x} \cdot \frac{(x-1) - (1+x)}{(x-1)^2}$$

$$= \frac{1}{2} \cdot \frac{-2}{(1+x)(x-1)} = \frac{1}{1-x^2}$$

(15)(i) Show that  $\operatorname{arcoth} x = \operatorname{artanh} \left( \frac{1}{x} \right)$

(ii) Find  $f(x)$  such that  $\operatorname{arcosh} x = \operatorname{arsinh}(f(x))$

### Solution

(i) Let  $y = \operatorname{arcoth} x$ , so that  $\operatorname{coth} y = x$

$$\Rightarrow \operatorname{tanh} y = \frac{1}{x}$$

$$\Rightarrow y = \operatorname{artanh} \left( \frac{1}{x} \right)$$

(ii) Let  $y = \operatorname{arcosh} x$ , so that  $\operatorname{cosh} y = x$

$$\Rightarrow \operatorname{sinh} y = \sqrt{x^2 - 1}$$

$$\Rightarrow y = \operatorname{arsinh}(\sqrt{x^2 - 1}) ; \text{ ie } f(x) = \sqrt{x^2 - 1}$$

(16) Given that  $\int \frac{1}{\sqrt{x^2 - a^2}} dx = \operatorname{arcosh} \left( \frac{x}{a} \right)$ , and that

$\operatorname{arcosh} x = \ln(x + \sqrt{x^2 - 1})$ , justify the writing of the integral as  $\ln(x + \sqrt{x^2 - a^2})$

### Solution

$$\operatorname{arcosh} \left( \frac{x}{a} \right) = \ln \left( \frac{x}{a} + \sqrt{\frac{x^2}{a^2} - 1} \right) = \ln \left( \frac{x + \sqrt{x^2 - a^2}}{a} \right)$$

$= \ln(x + \sqrt{x^2 - a^2}) - \ln a$ , which only differs from

$\ln(x + \sqrt{x^2 - a^2})$  by a constant

(17) Given that  $\operatorname{arcosh} x = \ln(x + \sqrt{x^2 - 1})$ , show that if  $\operatorname{cosh} a = b$  then  $a = \ln(b \pm \sqrt{b^2 - 1})$

**Solution**

$$\operatorname{cosh} a = b \Rightarrow a = \pm \operatorname{arcosh} b = \pm \ln(b + \sqrt{b^2 - 1})$$

$$\text{And } -\ln(b + \sqrt{b^2 - 1}) = \ln\left(\frac{1}{b + \sqrt{b^2 - 1}}\right) = \ln\left(\frac{b - \sqrt{b^2 - 1}}{b^2 - (b^2 - 1)}\right)$$

$$= \ln(b - \sqrt{b^2 - 1})$$


$$\text{so that } \pm \ln(b + \sqrt{b^2 - 1}) = \ln(b \pm \sqrt{b^2 - 1})$$

(18) Sketch the following:


(i)  $y = \operatorname{artanh} x$  (ii)  $y = \operatorname{coth} x$  (iii)  $y = \operatorname{arcoth} x$

**Solution**

(i)



(ii)



(iii)

