

Induction (3 pages; 2/6/23)

See STEP: "Misc. Topic Notes" re. Strong Induction

$$(1) \text{ Example: } \sum_{r=1}^n r^3 = \frac{1}{4}n^2(n+1)^2$$

Solution

$\sum_{r=1}^1 r^3 = 1^3 = 1$ and $\frac{1}{4}(1)^2(1+1)^2 = 1$; thus the result is true for $n = 1$

[Be careful to give enough working for both sides.]

Now assume that the result is true for $n = k$, so that

$$\sum_{r=1}^k r^3 = \frac{1}{4}k^2(k+1)^2$$

[A common error is to write "Let $n = k$ ", or "Assume $n = k$ ", or "If $n = k$ "]

[At this point it is possible to indicate the 'target' for $n = k + 1$; namely that $\sum_{r=1}^{k+1} r^3 = \frac{1}{4}(k+1)^2(k+2)^2$]

$$\begin{aligned} \text{Then } \sum_{r=1}^{k+1} r^3 &= \frac{1}{4}k^2(k+1)^2 + (k+1)^3 \\ &= \frac{1}{4}(k+1)^2(k^2 + 4(k+1)) \\ &= \frac{1}{4}(k+1)^2(k+2)^2 \\ &= \frac{1}{4}(k+1)^2([k+1] + 1)^2 \end{aligned}$$

But this is the result with k replaced by $k + 1$.

[Or, if the target has been mentioned previously, just indicate that the target has been obtained.]

So, if the result is true for $n = k$, then it is true for $n = k + 1$.

[A common error is to write: "So the result is true for $n = k$ and $n = k + 1$ "]

As the result is true for $n = 1$, it is therefore true for $n = 2, 3, \dots$ and, by [the principle of mathematical] induction, for all integer $n \geq 1$. [In some cases, it is appropriate to start at a different value for n , such as 0 or 2. This depends on what values of n the given formula is defined for.]

[Note that no credit is ever given in an exam for the standard wording on its own, or where the algebra is 'fudged'.]

(2) Example:

If $u_{n+1} = 2u_n + 3$, where $u_1 = 5$, then $u_n = 2^{n+2} - 3$

Solution

[Show that the result is true for $n = 1$]

Now assume that the result is true for $n = k$, so that

$$u_k = 2^{k+2} - 3$$

$$\begin{aligned} \text{Then } u_{k+1} &= 2u_k + 3 = 2(2^{k+2} - 3) + 3 = 2^{k+3} - 3 \\ &= 2^{(k+1)+2} - 3 \end{aligned}$$

[Standard wording]

(3) Example: $7^n + 4^n + 1$ is divisible by 6

Solution

[Show that the result is true for $n = 1$]

Now assume that the result is true for $n = k$

Approach 1

so that $7^k + 4^k + 1 = 6M$, where $M \in \mathbb{Z}^+$

To show that the result is then true for $n = k + 1$:

$$\begin{aligned} 7^{k+1} + 4^{k+1} + 1 &= 7(7^k + 4^k + 1) - 3(4^k) - 6 \\ &= 7(6M) - 6(2)(4^{k-1}) - 6 \\ &= 6(7M - 2(4^{k-1}) - 1), \text{ which is a multiple of 6 for } k \geq 1 \end{aligned}$$

(the multiple is positive, as $7^{k+1} + 4^{k+1} + 1$ is positive)

[Standard wording]

Approach 2

Let $f(k) = 7^k + 4^k + 1$

Then $f(k + 1) - \lambda f(k) = (7^{k+1} + 4^{k+1} + 1) - \lambda(7^k + 4^k + 1)$

[an appropriate λ will be chosen shortly]

$$= 7^k(7 - \lambda) + 4^k(4 - \lambda) + 1 - \lambda$$

Let $\lambda = 7$, so that $f(k + 1) - 7f(k) = -3(4^k) - 6$

and $f(k + 1) = 7f(k) - 6(2)(4^{k-1}) - 6$

As $f(k)$ is assumed to be a multiple of 6, and the other terms on the RHS are also multiples of 6 (for $k \geq 1$), it follows that

$f(k + 1)$ is a multiple of 6 (the multiple is positive, as

$7^{k+1} + 4^{k+1} + 1$ is positive).

[Standard wording]

[Textbooks sometimes consider $f(k + 1) - f(k)$ (ie with $\lambda = 1$), but this isn't guaranteed to work. (See Induction Exercises for examples of this.)]