Parabolas - Exercises (3 pages; 17/2/20)

See also the separate note "Parabolas" for further exercises.

Key to difficulty:

* easier


** moderate

*** harder

 (1^{***}) Using the parametric equations of a parabola $(x = at^2, y = 2at)$, show that the midpoints of chords of a parabola that have the same direction lie on a straight line parallel to the *x*-axis.

[A chord of a parabola joins two points on the parabola.]

 (2^{***}) A ray (eg of light) travels on a path parallel to the *x*-axis and hits the surface of the parabola $y^2 = 4ax$ at the point P $(at^2, 2at)$. The angle between the incoming ray and the normal at P is α . It can be assumed that the angle that the reflected ray makes with the normal is also α .



- (i) Show that $tan\alpha = t$
- (ii) Find the gradient of the reflected ray.

(iii) Show that the reflected ray passes through the focus of the parabola.

(3***) Suppose that $P(ap^2, 2ap)$ and $Q(aq^2, 2aq)$ are two points on the parabola $y^2 = 4ax$, such that the chord PQ passes through the focus of the parabola. Show that pq = -1.

(4***) If the tangents to a parabola at P and Q are perpendicular, show that the chord PQ passes through the focus S of the parabola.

(5***) Find the cartesian equations of the parabolas with:

(i) focus (4,4) and directrix y = 0

(ii) focus (2,2) and directrix x + y + 2 = 0

(6***) Suppose that $P(ap^2, 2ap)$ and $Q(aq^2, 2aq)$ are two points on the parabola $y^2 = 4ax$, such that the chord PQ passes through the focus of the parabola. Show that the tangents at P and Q meet on the directrix.

 $(7^{***}) P(ap^2, 2ap)$ and $Q(aq^2, 2aq)$ are two points on the parabola $y^2 = 4ax$, such that the chord PQ passes through the focus of the parabola. Show that the locus of the midpoint of PQ is a parabola, and establish its focus and directrix.