

Arithmetic Sequences & Series - Exercises (Solutions)

(6 pages; 22/3/20)

(1*) If teams A, B, C, D & E in some sporting competition have to play each other once, how many games are there in total?

Solution


AvB, AvC, AvD, AvE 4 games

BvC, BvD, BvE 3 games

CvD, CvE 2 games

DvE 1 game

Total = $1 + 2 + 3 + 4 = 10$ games



(2**) Extend (1) to find a formula for $1 + 2 + 3 + \dots + n$

Ideas


Consider the case $n = 4$

Use the diagram in (1)

Consider areas in the diagram

Give the areas letters

Can an equation be set up?



total of = X
 total of = Y
 total of = Z

Solution

Divide the 5×5 square up into the areas X, Y & Z

Let the squares be of unit area.

Then $X = 1 + 2 + 3 + 4$,

$Y = 5$ & $Z = X$

So, for $n = 4$, $X + 5 + X = 25$

Generalising this, $2X + (n + 1) = (n + 1)^2$

$$\Rightarrow 2X = (n + 1)[(n + 1) - 1] = (n + 1)n$$

$$\Rightarrow X = \frac{n(n+1)}{2}$$

(3*) For each of the following arithmetic sequences, find an expression for a_k :

(a) in the form $a_k = p + q(k - 1)$

(b) in the form $a_k = mk + c$

(c) in the form $a_k = a_{k-1} + t ; a_1 = u \quad (k \geq 2)$

(where p, q, m, c, t & u are to be found)

(i) 4, 7, 10, 13, 16, ...

Solution

(a) $a_k = 4 + 3(k - 1)$

(b) $a_k = 3k + 1$

(c) $a_k = a_{k-1} + 3 ; a_1 = 4 \quad (k \geq 2)$

(ii) -2, -1, 0, 1, 2, ...

Solution

(a) $a_k = -2 + (k - 1)$

(b) $a_k = k - 3$

(c) $a_k = a_{k-1} + 1 ; a_1 = -2 \quad (k \geq 2)$

(iii) 8, 6, 4, 2, 0, ...

Solution

(a) $a_k = 8 - 2(k - 1)$

(b) $a_k = 10 - 2k$

(c) $a_k = a_{k-1} - 2 ; a_1 = 8 \quad (k \geq 2)$

(4*) If $a_3 = 7$ and $a_{10} = 42$ are terms in an arithmetic sequence, find an expression for a_k .

Solution

Let $a_k = mk + c$

Then $7 = 3m + c$ & $42 = 10m + c$

Hence $35 = 7m$, so that $m = 5$ & $c = -8$

ie $a_k = 5k - 8$

(5**) Find

(i) $\sum_{k=1}^{20} (2k + 3)$

Solution

We know how to find $\sum_{k=1}^{20} k$

Consider $\sum_{k=1}^{20} 2k$

It equals $2 \sum_{k=1}^{20} k$

$$\sum_{k=1}^{20} (2k + 3) = \left(\sum_{k=1}^{20} 2k \right) + \left(\sum_{k=1}^{20} 3 \right)$$

$$\sum_{k=1}^{20} 3 = 3 + 3 + 3 + \dots + 3 = (20)(3)$$

So $\sum_{k=1}^{20} (2k + 3) = (2 \sum_{k=1}^{20} k) + (20)(3)$

$$= 2 \left(\frac{20(20+1)}{2} \right) + 60$$

$$= 420 + 60 = 480$$

$$(ii) \sum_{k=1}^{40} (10 - 4k)$$

Solution

$$\begin{aligned} \sum_{k=1}^{40} (10 - 4k) &= (10 \sum_{k=1}^{40} 1) - (4 \sum_{k=1}^{40} k) \\ &= 10(40) - 4 \left(\frac{40(40+1)}{2} \right) = 400 - 3280 = -2880 \end{aligned}$$

$$(6^{**}) \text{ Solve the equation } \sum_{k=1}^n (100 - 5k) = 0$$

Solution

$$\begin{aligned} \sum_{k=1}^n (100 - 5k) = 0 &\Rightarrow 100 \sum_{k=1}^n 1 = 5 \sum_{k=1}^n k \\ \Rightarrow 100n &= 5 \left(\frac{n(n+1)}{2} \right) \\ \Rightarrow 40n &= n(n+1) \Rightarrow n(n-39) = 0 \end{aligned}$$

Hence either $n = 0$ (not possible), or $n = 39$.

[Check: With $a = 100 - 5 = 95$, $d = -5$, $n = 39$,

$$\frac{n}{2} [2a + (n-1)d] = \frac{39}{2} [190 + 38(-5)] = 0]$$

$$(7^*) \text{ For each of the arithmetic sequences in (5), find } \sum_{k=1}^{100} a_k$$

Solution

$$(i) 4, 7, 10, 13, 16, \dots$$

$$\sum_{k=1}^{100} a_k = \frac{100}{2} (2(4) + 3(100 - 1)) = 15250$$

$$(ii) -2, -1, 0, 1, 2, \dots$$

$$\sum_{k=1}^{100} a_k = \frac{100}{2} (2(-2) + (100 - 1)) = 4750$$

(iii) 8, 6, 4, 2, 0, ...

$$\sum_{k=1}^{100} a_k = \frac{100}{2} (2(8) + (-2)(100 - 1)) = -9100$$

(8*) If I pay £50 into a bank account, then £60 a year later, followed by £70 the following year, and so on, increasing by £10 each year,

(i) How long will it take for the amount I pay in each year to reach £200?

Solution

$$50 + 10(k - 1) = 200 \Rightarrow 10k = 160 \Rightarrow k = 16$$

So, at the start of the 16th year I will be paying in £200.

(ii) How long will it take for the amount in the bank account to reach £1000?

Solution

$$\text{Consider } \frac{n}{2} [(2(50) + 10(n - 1))] = 1000$$

$$\Rightarrow n(90 + 10n) = 2000$$

$$\Rightarrow n^2 + 9n - 200 = 0$$

$$\Rightarrow n = \frac{-9 \pm \sqrt{81 + 800}}{2} = 10.3 \text{ (as } n > 0)$$

So, at the start of the 11th year (after paying in the amount due then) there will be over £1000 in the account.

(9**) For an arithmetic sequence with 1st term a and common difference d , show that the sum of the 1st n terms is

$$\frac{n}{2}[2a + (n - 1)d] \text{ by starting with } \sum_{k=1}^n [a + (k - 1)d]$$

Solution

$$\sum_{k=1}^n [a + (k - 1)d] = [(a - d) \sum_{k=1}^n 1] + d \sum_{k=1}^n k$$

$$= (a - d)n + d \cdot \frac{1}{2} n(n + 1)$$

$$= \frac{n}{2}(2a - 2d + dn + d) = \frac{n}{2}[2a + (n - 1)d]$$