

Counting (5 pages; 7/7/20)

Contents

(A) Permutations & Combinations

(B) Possible Approaches

(A) Permutations & Combinations

(1) Number of ways of arranging 5 items where order is important

There are 5 ways of filling the 1st place.

Then, for each of these, there are 4 ways of filling the 2nd place; then for each of these 20 ways, there are 3 ways of filling the 3rd place, and so on.

So the number of ways = $5 \times 4 \times 3 \times 2 \times 1 = 5!$

This is a special case of (2):

(2) Number of ways of selecting 2 items from 5 where order is important (“Permutations”)

eg placing 2 horses in a 5-horse race (P for place \Rightarrow P for Permutation)

Method 1

There are 5 ways of selecting the 1st item. Then, for each of these 5 ways, there are 4 ways of selecting the 2nd item.

Hence, ${}^5P_2 = 5 \times 4$

Method 2 (more complicated, but helps to understand Combinations - covered below)

From (1), the number of ways of arranging all 5 items is 5!

ABCDE, ABCED, ABDCE, ABDEC, ABECD & ABEDC all count as the same selection, if we are only interested in the first two letters – and there are 3! of these (the number of ways of arranging CDE). Similarly for all other pairs of 2 letters.

Therefore we need to divide by 3! to remove the duplication, to get:

$${}^5P_2 = \frac{5!}{3!} = 5 \times 4$$

This method of removing duplication will be used again below.

(3) Number of ways of selecting 2 items from 5 where order is NOT important (“Combinations”)

eg choosing 2 people out of 5 to form a Committee (C for Committee \Rightarrow C for Combination)

Because order is not important, ABCDE is treated as the same thing as BACDE.

So the 5P_2 ways in (2) contain duplication which is removed, as before, by dividing by 2! (the number of ways of arranging 2 letters)

$$\text{Hence } {}^5C_2 = \frac{5!}{3!2!}$$

(4) Number of ways of arranging 3 Ss and 2Fs, where the pattern of Ss and Fs is important

This can represent the possible arrangements of successes and failures in 5 trials, to give the coefficient in the Binomial probability.

Let the 3 Ss be labelled S_1 , S_2 & S_3 and the 2 Fs, F_1 and F_2 .

There are $5!$ ways of arranging the letters if they are thought of as all different.

However, if the Ss are considered to be indistinguishable, all the following count as the same arrangement:

$S_1F_1S_2S_3F_2$

$S_1F_1S_3S_2F_2$

$S_2F_1S_1S_3F_2$

$S_2F_1S_3S_1F_2$

$S_3F_1S_1S_2F_2$

$S_3F_1S_2S_1F_2$

As before, the duplication is removed by dividing by $3!$

If the Fs are also to be treated as indistinguishable, then we do the same thing with them and divide by $2!$

So the number of ways of arranging 3 Ss and 2Fs, where the pattern of Ss and Fs is important is: $\frac{5!}{3!2!}$

Although this is a case where order is important (in that the pattern of Ss and Fs is important), it has the same form as 5C_2 .

This can also be explained as follows:

We are interested in the different places that the Fs can occupy.

For example, SSSFF \Rightarrow places 4&5 SSFSF \Rightarrow places 3&5

As 'places 4&5' and 'places 5&4' would count as the same thing, the answer is 5C_2 (the number of ways of selecting 2 places from 5, where order doesn't matter).

(5) Interpretation of 5C_2 as the Binomial coefficient in the expansion of $(a+b)^5$

$$(a+b)^5 = (a+b)(a+b)(a+b)(a+b)(a+b)$$

The terms involving a^2 (and b^3) are obtained by selecting 2 of the 5 brackets (these 2 give rise to the “a”s and the other 3 give rise to the “b”s)

This can be done in 5C_2 ways [from (3)]; ie the term a^2b^3 occurs 5C_2 times.

(B) Possible Approaches

(a) Break down into various cases, and multiply by appropriate factors to take account of the number of possible arrangements for each case

(b) number of ways of allocating (eg) people to the 1st (eg) position \times number of ways of allocating people to the 2nd position etc

(c) Find total including non-permissible cases and then deduct these cases.

(d) If items have to be next to each other, combine them into a single block of r items, and multiply by $r!$

(e) For r indistinguishable items, assume initially that they are different, and then remove duplication by dividing by $r!$ (or vice-versa)

(f) Listing of possible situations

(g) Use of a Venn diagram to deal with overlapping cases

(h) For probabilities, either (1) $\frac{\text{number of favourable outcomes}}{\text{number of possible outcomes}}$

(assuming that the outcomes are equally likely), where combinations can often be used to find the numbers

or (2) $\text{Prob}(\text{eg } 1^{\text{st}} \text{ place being filled in a given way}) \times \text{Prob}(\text{2nd place being filled in a given way}) \times \dots$ (considering different cases separately, as in (a))