MAT Exercises – Transformations - Sol'ns

(9 pages; 4/11/22)

(1) Reflection in the line x = L: What composite transformation is equivalent to this?

translation of $\binom{-L}{0}$, followed by reflection in *y*-axis, followed by translation of $\binom{L}{0}$ $y = f(x) \rightarrow ?$ translation of $\binom{-L}{0}$: $y = f(x) \rightarrow y = f(x + L)$; reflection in *y*-axis: $y = f(x + L) \rightarrow y = f(-x + L)$ translation of $\binom{L}{0}$: $y = f(-x + L) \rightarrow y = f(-[x - L] + L)$ = f(2L - x) (2) What combination of transformations converts $y = 2^x$ to $y = 2^{4x-2}$?

 $y = 2^x \rightarrow y = 2^{4x}$ is a stretch of scale factor $\frac{1}{4}$ in the *x*-direction

Then
$$y = 2^{4x} \rightarrow y = 2^{4(x-\frac{1}{2})} = 2^{4x-2}$$
 is a translation of $\begin{pmatrix} \frac{1}{2} \\ 0 \end{pmatrix}$

[Alternatively, $y = 2^{4x} \rightarrow y = \left(\frac{1}{4}\right)2^{4x} = 2^{4x-2}$ is a stretch of scale factor $\frac{1}{4}$ in the *y*-direction.]

(3) (i) Find a series of transformations that can be applied to $y = \frac{1}{x}$ to produce $y = \frac{3x-2}{6x-1}$.

(ii) Hence or otherwise, sketch the curve $y = \frac{3x-2}{6x-1}$.

(i)
$$\frac{3x-2}{6x-1} = \frac{3x-\frac{1}{2}-\frac{3}{2}}{6x-1} = \frac{1}{2} - \frac{3}{12}\left(\frac{1}{x-\frac{1}{6}}\right)$$

So a possible series of transformations is:

a translation of
$$\begin{pmatrix} \frac{1}{6} \\ 0 \end{pmatrix}$$
,

followed by a stretch of scale factor $\frac{1}{4}$ in the *y*-direction,

followed by a reflection in the *x*-axis,

followed by a translation of $\begin{pmatrix} 0\\ \frac{1}{2} \end{pmatrix}$

[Note: $\frac{1}{2} - \frac{3}{12} \left(\frac{1}{x - \frac{1}{6}} \right) = \frac{1}{2} - \frac{1}{4x - \frac{2}{3}}$, so an alternative series of transformations is:

a translation of $\begin{pmatrix} \frac{2}{3} \\ 0 \end{pmatrix} \begin{bmatrix} \frac{1}{x} \to \frac{1}{x - \frac{2}{3}} \end{bmatrix}$ followed by a stretch of scale factor $\frac{1}{4}$ in the *x*-direction $\begin{bmatrix} \frac{1}{x - \frac{2}{3}} \to \frac{1}{4x - \frac{2}{3}} \end{bmatrix}$, followed by a reflection in the *x*-axis, followed by a translation of $\begin{pmatrix} 0 \\ \frac{1}{2} \end{pmatrix}$.


Alternatively, $\frac{1}{4x-\frac{2}{3}}$ could be obtained instead by a stretch of scale factor $\frac{1}{4}$ in the *x*-direction $[\frac{1}{x} \rightarrow \frac{1}{4x}]$ (or a stretch of scale factor $\frac{1}{4}$ in the *y*-direction

$$\left[\frac{1}{x} \to \frac{1}{4}\left(\frac{1}{x}\right)\right]$$
, followed by a translation of $\begin{pmatrix}\frac{1}{6}\\0\end{pmatrix}\left[\frac{1}{4x} \to \frac{1}{4\left(x-\frac{1}{6}\right)}\right]$.

fmng.uk

(ii) As an alternative to performing the transformations in (i):

Step 1: $x = 0 \Rightarrow y = 2$; $y = 0 \Rightarrow x = \frac{2}{3}$ Step 2: vertical asymptote when $6x - 1 = 0 \Rightarrow x = \frac{1}{6}$ $x = \frac{1}{6} + \delta$ ($\delta > 0$ is small) $\Rightarrow y = \frac{3x-2}{6x-1} = \frac{-}{+}$; ie y < 0 $[x = \frac{1}{6} - \delta \Rightarrow y = \frac{-}{-}$; ie y > 0] Step 3: $\lim_{x \to \infty} \frac{3x-2}{6x-1} = \lim_{x \to \infty} \frac{3-\frac{2}{x}}{6-\frac{1}{x}} = \frac{3}{6} = \frac{1}{2}$ (and also as $x \to -\infty$) Step 4: When x = 100, $y = \frac{298}{599} < \frac{1}{2}$, so that $y \to \frac{1}{2}^{-}$ as $x \to \infty$ and when x = -100, $y = \frac{-302}{-601} > \frac{1}{2}$, so that $y \to \frac{1}{2}^{+}$ as $x \to -\infty$



(4) What combination of transformations converts $y = 3^{-x}$ to $y = 3^{2x-1}$?

 $y = 3^{-x} \rightarrow y = 3^x$ is a reflection in the *y*-axis.

 $y = 3^x \rightarrow y = 3^{x-1}$ is a translation of $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$

 $y = 3^{x-1} \rightarrow y = 3^{2x-1}$ is a stretch of scale factor $\frac{1}{2}$ in the

x-direction