


## STEP 2007, Paper 2, Q2 – Solution (3 pages; 23/5/18)



(i)  $y = 2x^3 - bx^2 + cx$

$$\frac{dy}{dx} = 6x^2 - 2bx + c$$

Turning point  $\Rightarrow 6x^2 - 2bx + c = 0$  (1)

The roots of (1) are  $p$  &  $q$ ,

so that  $p + q = \frac{2b}{6}$  &  $pq = \frac{c}{6}$

Hence  $b = 3(p + q)$  &  $c = 6pq$

(ii) The point of inflexion is the turning point of the gradient;

ie where  $\frac{d}{dx} \left( \frac{dy}{dx} \right) = 0$

So  $\frac{d^2y}{dx^2} = 12x - 2b = 0$ , and hence  $x = \frac{b}{6} = \frac{p+q}{2}$

ie the point of inflexion lies halfway between the turning points.

There is rotational symmetry (of order 2) about the point of inflexion.

[All cubics have a single point of inflexion, and it always lies halfway between the turning points (if they exist). It isn't clear whether a proof of the symmetry is required for this question. It would be strange for no justification to be needed, but I would be surprised if the following was intended. See "Cubic Functions" for an alternative method.]

To demonstrate rotational symmetry about  $x = \frac{b}{6}$ , we require that

$$f\left(\frac{b}{6} + t\right) - f\left(\frac{b}{6}\right) = -\left\{f\left(\frac{b}{6} - t\right) - f\left(\frac{b}{6}\right)\right\};$$

$$\text{ie that } f\left(\frac{b}{6} + t\right) + f\left(\frac{b}{6} - t\right) = 2f\left(\frac{b}{6}\right)$$

$$\text{where } f(x) = 2x^3 - bx^2 + cx$$

$$\text{Now } f\left(\frac{b}{6}\right) = \frac{2b^3}{6^3} - \frac{b^3}{36} + \frac{bc}{6}$$

$$\text{and } f\left(\frac{b}{6} + t\right) + f\left(\frac{b}{6} - t\right) = 2\left(\frac{b}{6} + t\right)^3 - b\left(\frac{b}{6} + t\right)^2 + c\left(\frac{b}{6} + t\right)$$

$$+ 2\left(\frac{b}{6} - t\right)^3 - b\left(\frac{b}{6} - t\right)^2 + c\left(\frac{b}{6} - t\right)$$

$$= \frac{4b^3}{6^3} + 4(3)\left(\frac{b}{6}\right)t^2 - \frac{2b^3}{36} - 2bt^2 + \frac{2bc}{6}$$


$$= \frac{4b^3}{6^3} - \frac{2b^3}{36} + \frac{2bc}{6} = 2f\left(\frac{b}{6}\right), \text{ as required}$$

$$\text{(iii) } m - n = f(p) - f(q)$$

$$= 2(p^3 - q^3) - b(p^2 - q^2) + c(p - q)$$

$$\begin{aligned}
&= 2(p - q)(p^2 + pq + q^2) - 3(p + q)(p^2 - q^2) + 6pq(p - q) \\
&= (p - q)\{2p^2 + 2pq + 2q^2 - 3(p + q)^2 + 6pq\} \\
&= (p - q)\{2(p + q)^2 - 2pq - 3(p + q)^2 + 6pq\} \\
&= (p - q)\{4pq - (p + q)^2\} \\
&= -(p - q)(p - q)^2 \\
&= (q - p)^3, \text{ as required}
\end{aligned}$$

(iv)



$R = 1/2 \times$  area of rectangle in diagram above (by symmetry of the cubic about the point of inflexion)

$$\begin{aligned}
&= \frac{1}{2}(q - p)(m - n) \\
&= \frac{1}{2}(q - p)(q - p)^3 \\
&= \frac{1}{2}(q - p)^4, \text{ as required}
\end{aligned}$$